



10 / 524085
PCT/GB 2003 / 003447



INVESTOR IN PEOPLE

19 FEB 2003

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

RECD 04 SEP 2003

WIPO PCT

PRIORITY DOCUMENT


SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed *Stephen Hordley*

Dated 22 August 2003

BEST AVAILABLE COPY

The
Patent
Office

THE PATENT OFFICE
D

14 AUG 2002

NEWPORT

1/77

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

1. Your reference

11101P6 GB/AB

2. Patent application number

(The Patent Office will fill in this part)

0218857.1

14AUG02 E740832-1 D02903

P01/7700 0.00-0218857.1

14 AUG 2002

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Reckitt Benckiser N.V.
Kantoorgebouw De Appelaer
De Fruittuinen 2-12
2132 NZ Hoofddorp
NETHERLANDS

Patents ADP number (if you know it)

07921075005 ✓

If the applicant is a corporate body, give the country/state of its incorporation

Netherlands

4. Title of the invention

Bactericide Surfactant Compositions

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Andrew S Brown
Reckitt Benckiser plc
Group Patents Department
Dansom Lane
HULL
HU8 7DS
UNITED KINGDOM

Patents ADP number (if you know it)

07799521001 ✓

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application
(day / month / year)

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

Yes

a) any applicant named in part 3 is not an inventor, or

b) there is an inventor who is not named as an applicant, or

c) any named applicant is a corporate body.

See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description	27
Claim(s)	3
Abstract	--
Drawing(s)	--

10. If you are also filing any of the following, state how many against each item.

Priority documents
Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

One ✓

Request for substantive examination (Patents Form 10/77)

One /

Any other documents (please specify)

FS2

11. I/We request the grant of a patent on the basis of this application.

Signature

Date

John C McKnight

John C McKnight

9 August 2002

12. Name and daytime telephone number of Person to contact in the United Kingdom

Andrew S Brown (01482) 582411

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- i) If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- ii) Write your answers in capital letters using black ink or you may type them.
- iii) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- iv) If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- v) Once you have filled in the form you must remember to sign and date it.
- vi) For details of the fee and ways to pay please contact the Patents Office.

BACTERICIDE SURFACTANT COMPOSITIONS

The invention relates to anionic surfactant compositions containing a bactericide, which is a cationic compound, being specifically developed for fabric cleaning products.

In general, cationic bactericide compounds have low compatibility with anionic based surfactant compositions and they may have a negative effect in terms of the cleaning performance of the composition.

Certain cationic compounds, as described below, are well known actives able to provide good disinfecting efficacy both versus gram negative and gram positive bacteria, but have in general low compatibility with anionic surfactants or anionic species in general. Cationic compounds tend to precipitate with anionic compounds or lose their bactericide efficacy due to the formation of an anionic-cationic complex that doesn't allow the cationic part of the complex to be available to kill bacteria. The cationic compounds are, in addition, well known to cause stickiness to fabric surfaces, especially carpet, and are very deleterious for soil re-deposition, which is considered an important factor for fabric care. Generally cationic compounds are metal corrosives making their use in aerosols problematical.

Phenols and phenol based bactericides are other ingredients used in several application areas. More and more concerns about their toxicological impact is growing, they are considered as potential carcinogens and they are generally avoided for this reason.

Aldehydes such as formaldehyde and glutaraldehyde are cheap and broad-spectrum bactericides, but as with phenols, they are carcinogens or potential carcinogens and also show a tendency to sensitise people who are frequently exposed to them.

The halogens have a long history as bactericide agents. Chloride is the active atom in household bleach and chloride and bromine are used for water disinfection. Iodine is

very commonly incorporated into antiseptics, for disinfecting skin and wound dressings, and it is added to water for water treatment. Iodine used as such or in combination with organic carrier molecules, iodophors, is used as a liquid disinfectant, but it tends to stains the treated surface with reddish-brown colour.

5 Chlorine is cheap and very effective, but it tends to corrode metal surface and to decolourise dyes from the fabric surface.


Organic acids are known in the art to be bactericides, examples are citric acid and salicylic acid. The organic acids are efficient at low pH, below 5 and more preferably
10 below 4. At these low pH conditions the cleaning performances of the surfactants are reduced and the compositions work mainly as a bactericide not as a good cleaning product.

Alcohols, such as iso-propanol and ethanol, have been used for a long time in ready
15 to use disinfectants for medical and consumer products. To be effective they need to make up a significant percentage of the composition, usually 15-70 % w/v, the disadvantages are that, being a solvent, they attack many polymers and plastics and cause the composition to be high flammable.

20 Other known ingredients are the essential oils, such as tea tree oil, thyme oil and citronella oil. These products show a low/ medium bacteria efficacy, unless used at high concentration, but at high concentrations they may become sensitisers.

In the present invention we have surprisingly found that certain cationic compounds,
25 as described below, are compatible with anionic surfactants and anionic based products and maintain their bactericidal activity, having a low toxicological impact and low or no negative effects on the composition in terms of cleaning, soil re-deposition and fibre damage.

30 The cationic compounds of the invention are those that provide a germicidal effect to the concentrate compositions, and especially preferred are quaternary ammonium salts which may be characterised by the general structural formula:




5

wherein R1, R2, R3 and R4 are independently selected from alkyl, aryl or alkylaryl substituent of from 1 to 26 carbon atoms, and the entire cation portion of the molecule has a molecular weight of at least 165. The alkyl substituents may be long-chain alkyl, long-chain alkoxyaryl, long-chain alkylaryl, halogen-substituted long-chain alkylaryl, long-chain alkylphenoxyalkyl and arylalkyl. The remaining substituents on the nitrogen atoms other than the above mentioned alkyl substituents are hydrocarbons usually containing no more than 12 carbon atoms. The substituents R1, R2, R3 and R4 may be straight-chained or may be branched, but are preferably straight-chained, and may include one or more amide, ether or ester linkages.

15

Preferred cationic compounds of the invention which are useful in the practice of the present invention include those which have the structural formula:

20



25


wherein R2 and R3 are each independently the same or different C8 -C12 alkyl; or R2 is selected from C12-16 alkyl, C8-18 alkylethoxy or C8-18 alkylphenoxyethoxy and R3 is benzyl. Counterion X⁻ is a salt forming anion as described below. The alkyl groups recited in R2 and R3 may be straight-chained or branched, but are preferably substantially linear. Such useful quaternary compounds are available and include ONYXIDE™ 3300 is described as n-alkyl dimethyl benzyl ammonium

30

ethoxylate with 7 moles EO; Tomadol 45-7, a C14-C15 linear primary alcohol ethoxylate with 7 moles EO; and Tomadol 91-6, a C9-C11 linear alcohol ethoxylate with 6 moles EO.


- 5 A preferred surfactant is an anionic surfactant. Such anionic surface active agents are frequently provided in a salt form, such as alkali metal salts, ammonium salts, amine salts, amino alcohol salts or magnesium salts. Contemplated as useful are one or more sulfate or sulfonate compounds including: alkyl sulfates, alkyl ether sulfates, alkylamidoether sulfates, alkyl benzene sulfates, alkyl benzene sulfonates, alkylaryl
- 10 polyether sulfates, monoglyceride sulfates, alkylsulfonates, alkylamide sulfonates, alkylarylsulfonates, olefinsulfonates, paraffin sulfonates, alkyl sulfosuccinates, alkyl ether sulfosuccinates, alkylamide sulfosuccinates, alkyl sulfosuccinamate, alkyl sulfoacetates, alkyl carboxylates, alkyl phosphates, alkyl ether phosphates, acyl sarconsinates, acyl isethionates, and N-acyl taurates. Generally, the alkyl or acyl
- 15 radical in these various compounds comprise a carbon chain containing 12 to 20 carbon atoms.

Preferred surfactants are also alkyl naphthalene sulfonate anionic surfactants of the formula:



- wherein R is a straight chain or branched alkyl chain having from about 1 to about 25 carbon atoms, saturated or unsaturated, and the longest linear portion of the alkyl
- 30 chain is 15 carbon atoms or less on the average, M is a cation which makes the compound water soluble especially an alkali metal such as sodium or magnesium, ammonium or substituted ammonium cation.

Particularly preferred are alkyl sarcosinate, sulfosuccinate and alkyl sulfate anionic surfactants of the formula




wherein R is a straight chain or branched alkyl chain having from about 8 to about 18 carbon atoms, saturated or unsaturated, and the longest linear portion of the alkyl chain is 15 carbon atoms or less on the average, M is a cation which makes the compound water soluble especially an alkali metal such as sodium or magnesium, ammonium or substituted ammonium cation, and x is from 0 to about 4. Most preferred are the non-ethoxylated C12-15 primary and secondary alkyl sulfates, especially sodium lauryl sulfate.

Most desirably, the anionic surfactant according to constituent is selected to be of a type that dries to a friable powder. This facilitates their removal from carpets and carpet fibres, such as by brushing or vacuuming.



Super wetting agents are used between 0.01 to 10% w/v, preferably from 0.1 to 5% w/v. The super wetting agents of this invention are silicone glycol copolymers and fluorsurfactants.

The silicone glycol copolymers are described by the following formula:




Where x, y, m and n are whole number ranging from 0 to 25. X is preferred between 0-10 and y, m and n between 0-5. R and R' are straight chain or branched alkyl chain having from about 1 to about 25 carbon atoms, saturated or unsaturated, and the longest linear portion of the alkyl chain is 15 carbon atoms or less on the average.

5 The fluorinated surfactant is described in the following formulae:



10



Wherein n, m and x are integers having a value from 0 to 15; preferred values are between 1 and 12. M is a cation which makes the compound water-soluble especially an alkali metal such as sodium or magnesium, ammonium or substituted ammonium
15 cation.

The super wetting agents described are able to low the surface tension in water at values below 25 mN/m, in the range between 18 and 25 mN/m at concentrations of 0.0001 to 1% w/v, preferably between 0.001 to 0.1% w/v.

20

The composition of the present invention may also contain one or more hydrotropes. Examples of suitable hydrotropes are sodium cumene sulfonate (ELTESOL SC40 available from Albright & Wilson), sodium xylene sulfonate (ELTESOL SX40 available from Albright & Wilson), di-sodium mono- and di-alkyl disulfonate
25 diphenyloxide (DOWFAX 3B2 available from Dow Chemicals), n-octane sodium sulfonate (BIOTERGE PAS 7 S or 8 S available from Stepan). Levels of hydrotrope added are from 0.01% to 15% w/v.

Organic solvents may be added and may be beneficial in term of improving the
30 solubility of the cationic compounds in water. The organic solvents should be water-miscible. Preferably the organic solvent is found at levels of 0.01 to 15% w/v. The organic solvent constituent of the inventive compositions include one or more

alcohols, glycols, acetates, ether acetates and glycol ethers. Exemplary alcohols useful in the compositions of the invention include C2-C8 primary and secondary alcohols which may be straight chained or branched. Exemplary alcohols include pentanol and hexanol. Exemplary glycol ethers include those glycol ethers having
5 the general structure Ra-O-Rb-OH, wherein Ra is an alkoxy of 1 to 20 carbon atoms, or aryloxy of at least 6 carbon atoms, and Rb is an ether condensate of propylene glycol and/or ethylene glycol having from 1 to 10 glycol monomer units. Preferred are glycol ethers having 1 to 5 glycol monomer units.

10 By way of further non-limiting example specific organic constituents include propylene glycol methyl ether, dipropylene glycol methyl ether, tripropylene glycol methyl ether, propylene glycol n-propyl ether, ethylene glycol n-butyl ether, diethylene glycol n-butyl ether, diethylene glycol methyl ether, propylene glycol, ethylene glycol, isopropanol, ethanol, methanol, diethylene glycol monoethyl ether
15 acetate and particularly useful is , propylene glycol phenyl ether, ethylene glycol hexyl ether, diethylene glycol hexyl ether.

The chelating agent is added at a level between 0.01 to 5 % w/v, preferably between 0.1 to 1% w/v. Examples of chelating agents are described below:

20

- the parent acids of the monomeric or oligomeric polycarboxylate chelating agents or mixtures therefore with their salts, e.g. citric acid or citrate/citric acid mixtures are also contemplated as useful builder components.

25

- borate builders, as well as builders containing borate-forming materials than can produce borate under detergent storage or wash conditions can also be.

- iminosuccinic acid metal salts

30

- polyaspartic acid metal salts.

- examples of bicarbonate and carbonate builders are the alkaline earth and the alkali metal carbonates, including sodium carbonate and sesquicarbonate and mixtures thereof. Other examples of carbonate type builders are the metal carboxy glycine and metal glycine carbonate.

5

- ethylene diamino tetra acetic acid and salt forms.

- water-soluble phosphonate and phosphate builders are useful for this invention.

10

Examples of phosphate builders are the alkali metal tripolyphosphates, sodium potassium and ammonium pyrophosphate, sodium and potassium and ammonium pyrophosphate, sodium and potassium orthophosphate sodium polymeta/phosphate in which the degree of polymerisation ranges from 6 to 21, and salts of phytic acid.

15

Specific examples of water-soluble phosphate builders are the alkali metal tripolyphosphates, sodium potassium and ammonium pyrophosphate, sodium and potassium and ammonium pyrophosphate, sodium and potassium orthophosphate, sodium polymeta/phosphate in which the degree of polymerisation ranges from 6 to 21, and salts of phytic acid.

20

The polymers used in this invention at a level between 0.01 to 30% w/v, preferably between 0.1 to 5% w/v. Examples of polymers are: water-soluble compounds include the water soluble monomeric polycarboxylates, or their acid forms, homo or copolymeric polycarboxylic acids or their salts in which the polycarboxylic acid comprises at least two caroxylic radicals separated from each other by not more than two carbon atoms, carbonates, bicarbonates, borates, phosphates, and mixtures of any of thereof. The carboxylate or polycarboxylate builder can be monomeric or oligomeric in type although monomeric polycarboxylates are generally preferred for reasons of cost and performance. Suitable carboxylates containing one carboxy group include the water soluble salts of lactic acid, glycolic acid and ether derivatives thereof. Polycarboxylates containing two carboxy groups include the water-soluble salts of succinic acid, malonic acid, (ethylenedioxy) diacetic acid, maleic acid,

30

diglycolic acid, tartaric acid, tartronic acid and fumaric acid, as well as the ether carboxylates and the sulfinyl carboxylates. Polycarboxylates containing three carboxy groups include, in particular, water-soluble citrates, aconitrates and citraconates as well as succinate derivatives such as the carboxymethloxysuccinates
5 described in GB-A-1,379,241, lactoxysuccinates described in GB-A-1,389,732, and aminosuccinates described in NL-A-7205873, and the oxypolycarboxylate materials such as 2-oxa-1,1,3-propane tricarboxylates described in GB-A-1,387,447.

Polycarboxylate containing four carboxy groups include oxydisuccinates disclosed in
10 GB-A-1,261,829, 1,1,2,2-ethane tetracarboxylates, 1,1,3,3-propane tetracarboxylates and 1,1,2,3-propane tetracarboxylates. Polycarboxylates containing sulfo substituents include the sulfosuccinate derivatives disclosed in GB-A-1,398,421, GB-A-1,398,422 and US-A-3,936,448, and the sulfonated pyrrolidone citrates described in GB-A-1,439,000.

15

Alicyclic and heterocyclic polycarboxylates include cyclopentane-cis,cis,cis-tetracarboxylates, cyclopentadienide pentacarboxylates, 2,3,4,5,6-hexane – hexacarboxylates and carboxymethyl derivatives of polyhydric alcohols such as sorbitol, mannitol and xylitol. Aromatic polycarboxylates include mellitic acid,
20 pyromellitic acid and the phthalic acid derivatives disclosed in GB-A-1,425,343.

Of the above, the preferred polycarboxylates are hydroxycarboxylates containing up to three carboxy groups per molecule, more particularly citrates.

25 More preferred polymers are homo-polymers, copolymers and multiple polymers of acrylic, fluorinated acrylic, sulfonated styrene, maleic anhydride, metacrylic, isobutylene, styrene and ester monomers.

Examples of these polymers are Acusol supplied from Rohm & Haas, Syntran
30 supplied from Interpolymer and Versa and Alcosperse series supplied from Alco Chemical, a National Starch & Chemical Company.

Antifoaming agents are critical for machine shampoo products: they are used in this invention at a level between 0.01 to 5% w/v. The foam level in fact doesn't allow to properly use the carpet cleaning machines if it is too high and in any case tends to reduce the mechanical action of the carpet cleaner machine brushes, having an impact on soil removal. Atifoaming agents are so considered important actives of this invention.

Examples are polydimethylsiloxanes in combination with hydrophobic silica in different ratios.

Water is present in the compositions at levels of up to 98% w/v, ideally up to 90% w/v.

Points of advantage found in this invention:

- The cationic compounds of the invention are compatible with anionic surfactants and other anionic species such as anionic polymers.
- The cationic compounds of the invention, even if complexed provide a bactericidal action similar to uncomplexed cationic species.
- The cationic compounds in fabric cleaning compositions don't cause any reduction in terms of cleaning performance, fabric damage and anti- re-deposition.
- The cationic compounds can be combined with small amount of other bactericidal actives, such as essential oils, phenols, alcohols and acids, improving the bactericidal effect without lowering cleaning performance.

These cationic compounds can be used in liquid and powder carpet cleaning compositions. Examples of liquid compositions are ready to use products as triggers and dilutable products as manual and machine shampoos.

These cationic complexes are not very soluble in water, but this point can be easily overcome by combining them with anionic and non-ionic surfactants, by using solvents, hydrotropes and polymers. Heating the liquid compositions up to 60-80°C can also help to improve the dissolution and manufacture.

5

TESTS

Several tests have been considered in this invention to evaluate the compatibility of the cationic complexes in anionic carpet cleaning compositions (Storage stability test), the bactericidal efficacy of the cationic complexes (AOAC Germicidal and Detergent Sanitizing Action of Disinfectants method), the cleaning performances (Stain removal test), the anti re-deposition and the carpet damage.

10

The cleaning performances have been compared with Resolve spot & stain carpet cleaner trigger and with Resolve pet stain & odour carpet cleaner trigger.

15

STORAGE STABILITY TEST:

The 100 ml of the liquid compositions were put in to glass jars at 5, 25 and 40 °C. The formula appearance and pH variation are considered and a rating assigned. The storage situation is monitored after 1 month and 3 months and compared with the starting values. The data table with the corresponding storage rating is reported below:

20

Storage rating (SR)	pH variation	Liquid appearance
0	0	No difference from reference
1	Less than 0.25	No difference from reference
2	From 0.25 to 0.50	No difference from reference

3	From 0.5 to 1	No difference from reference
4	More than 1	No difference from reference
5	From 0.25 to 0.50	Separation and/or colour change
6	From 0.25 to 0.5	Separation and/or colour change
7	From 0.5 to 1	Separation and/or colour change
8	More than 1	Separation and/or colour change

The higher the rating number, the worse the storage performance. The anionic species found compatible with the cationic complexes are also reported.

5

BACTERICIDAL EFFICACY TEST:

The AOAC Germicidal and Detergent Sanitizing Action of Disinfectants method has been used for bactericidal efficacy evaluation.

- 10 The efficacy has been tested against *Staphylococcus aureus* (ATCC 6538) and *Enterobacter aerogenes* (ATCC13048) and it is expressed as percentage bacteria reduction.

STAIN REMOVAL TEST:

15

This method has been designed for the evaluation of stain removal performance of carpet cleaner compositions and especially of trigger products. The Resolve™ spot & stain carpet cleaner trigger and Resolve™ pet stain & odour carpet cleaner trigger has been used as a reference for spot removal products.

The stains were applied to nylon carpets following the procedures below:

- | | | |
|----|-------------------|--|
| 5 | - Dirty Motor Oil | 0.1 grams sprayed with an airbrush using a 5 cm diameter template. Allow any sediment to settle and decant oil into a clean container. |
| | - Ink | 0.1 grams sprayed with an airbrush using a 5 cm diameter template – Blue oily ink (Pelikan). |
| 10 | - Grape juice | 1.5 gram sprayed with an airbrush using a 5 cm diameter template – Welch's |
| | - Mud | 1.1 grams diluted 1:4 with water, well mixed and applied with a spatula into a loom with 3 cm diameter – green mud argital. |
| 15 | | 5 cm diameter template – 7 Lipton yellow teabags are put in 200 grams boiling water. |
| | - Coffee | 1 gram sprayed with an airbrush using a 5 cm diameter template – milk (3 ml) & sugar (20 g) |
| | - Spaghetti sauce | 0.75 grams applied with a spatula into a loom with 3 cm diameter – Ragù traditional. |

20

The stains were allowed to set for 24 hours before cleaning. The cleaning process was carried out by spraying 4 grams of the product onto the stain, leaving it for 3 minutes and doing 10 strokes in the vertical and then in the horizontal direction using a clean cloth. The area is then blotted twice and left to dry for 24 hours.

25

A panel test has been done on stained and cleaned carpet swatches to check the stain removal from carpet pile. The panellists are provided with a new carpet as a 1 reference and a stained carpet as a reference of 5. The panellists assigned a value to the cleaned carpet swatches between 1 and 5. The lower the rating number, the better the cleaning performance. The rating number obtained from carpet pile and backing per stain/product is averaged.

CARPET DAMAGE TEST:

Multiple (x3) cleaning process steps were done on a blue nylon carpet and red wool carpet to check for any potential damage of the compositions of this invention to the carpet fibre or colour. Resolve™ trigger spot & stain carpet cleaner was used as a reference product.

A panel test is used for evaluating carpet damage. Panellists are given a non-cleaned carpet swatch as a control of 0 per each carpet type/colour. The evaluation scale ranges from 0 to 4, 0 is no colour change, 4 corresponds to very high colour change. The responses of the panellist are averaged.

Damage rating (DR)	Carpet damage description
0	No colour change or fibre damage
1	Slight colour change and fibre damage
2	Perceptible colour change and fibre damage
3	Evident colour change and fibre damage
4	Strong colour change and fibre damage

An accepted carpet damage rating is parity or below 1.

15 **ANTI RE-DEPOSITION TEST:**

This method has been designed for the evaluation of anti re-deposition product evaluation.

The soil used has the following composition:

	Soil components:	% by weight
5	Peat Moss	47.7
	Cement	21.4
	Kaolin clay	8.0
	Silica	8.0
	Red Iron oxide	1.3
10	Charcoal	12.6
	Mineral oil	1.0

A nylon carpet is used for the test.

15 A cleaning process is carried out by spraying 4 grams of the product onto the carpet, leaving it for 3 minutes and doing 10 strokes in the vertical and then in the horizontal direction using a clean cloth. The area is then blotted twice and left to dry for 24 hours.

20 The carpet is soiled with 5 grams of standard soil. The soil is applied 1 gram once by strainer. The soiled carpet is then put with 4 kg of steel beads in the jar mill and stirred for 30 minutes at 56 rpm.

Carpet swatches are placed in a dark room temperature chamber (25°C / 50%RH) for
25 24 hours while they dry.

The carpet swatches are the vacuumed doing 3 strokes in the vertical and then in the horizontal direction.

30 The water is used as a 'product' reference for having no positive or negative impact on anti re-deposition.

The anti re-deposition performance is evaluated by measuring the carpet with a portable spectrophotometer before cleaning after soiling and vacuuming. The result is reported as anti re-deposition percentage, where a positive value indicates an anti re-deposition effect and a negative value indicates a product stickness.

5

EXAMPLES:

The liquid compositions are typically prepared by mixing each ingredient
10 together in a suitable container.

Examples of compositions forming a part of the present invention related to ready to use products are set below in Tab 1 and 2, examples for dilutable machine carpet cleaner liquids are reported in table 3 and examples for dilutable manual
15 shampoo liquids with are described in table 4. All the various components are identified in Tab 5.

Table 1					
Components	Ex 1	Ex 2	Ex 3	Ex 4	Ex 5
	%	%	%	%	%
Trilon B	11.20	11.20	11.20	1.00	1.00
Trilon BS acid	2.00	1.40	2.00		
Citric acid				0.11	0.08
Stepanol WAC	3.00	3.00	3.00	3.00	3.00
Syntran 1575	3.00	3.00	3.00		
Zelan 338				0.15	0.15
Genapol 26-L-5	0.50	0.50	0.50		
Zonyl 7950				0.15	0.15
Onyxide 3300 33%	0.50	0.30	0.30	0.35	0.35

Kathon CG/ICP	0.10	0.10	0.10	0.05	0.05
Thymol Kristal powder	0.09	0.09	0.09	0.09	0.09
Dowanol PPH		1.50	1.50	1.50	1.50
Isopropyl alcohol				2.00	2.00
Soft water	79.61	78.91	78.31	91.65	91.68
pH	6.7	8.0	6.5	6.5	8.0

	Table 2				
Components	Ex 6	Ex 7	Ex 8	Ex 9	Ex 10
	%	%	%	%	%
Trilon B	1.00	1.00	1.00	1.00	1.00
Citric acid	0.08	0.08	0.08	0.08	0.08
Stepanol WAC	3.00	3.00	3.00	3.00	3.00
Zelan 338	0.15	0.15	0.15	0.15	0.15
Zonyl 7950	0.15	0.15	0.15	0.15	0.15
Onyxdex 3300 33%	0.20	0.50	0.20	0.35	0.50
Kathon CG/ICP	0.05	0.05	0.05	0.05	0.05
Thymol Kristal powder	0.09	0.09			
Fragrance			0.15	0.15	0.15
Dowanol PPH	1.50	1.50	1.50	1.50	1.50
Isopropyl alcohol	2.00	2.00	2.00	2.00	2.00
Soft water	91.78	91.48	91.72	91.57	91.42
pH	7.8	7.8	7.7	7.7	7.7

	Table 3		
Components	Ex 11	Ex 12	Ex 13
	%	%	%
Trilon B	8.11	8.11	8.11
Trilon BS acid	0.77	0.77	0.77
Sodium bicarbonate	0.06	0.06	0.06
Petro 11 liquid	1.23	1.23	1.23
Eltesol SC40	7.00	6.00	9.50
Genapol 26-L-60	2.03	2.03	2.03
Syntran 1580	1.59	1.59	1.59
Onyxide 3300	1.90	1.90	1.90
Silicone antifoam	0.05	0.05	0.05
Fragrance	0.40	0.40	0.40
Dowanol PPH			1.00
Dowanol DPnP	1.90	2.85	
Soft water	74.96	75.01	73.362
pH	8.7	8.9	8.7

	Table 4	
Components	Ex 14	Ex 15
	%	%
Dequest 2066	2.00	2.00
Empicol 0335	19.00	19.00
Syntran DX 302-1	12.50	12.50
Onyxide 3300 33%	1.40	2.80
Fragrance	0.25	0.25
Dye	0.0004	0.0004
Soft water	64.85	63.45
pH	6.9	6.9

	Table 5
Component	Description of component
Trilon B	EDTA tetrasodium salt 40% from BASF
Trilon BS acid	EDTA acid powder from BASF
Citric acid	Citric acid from various suppliers
Sodium bicarbonate	Sodium bicarbonate from various suppliers
Syntran 1575	Fluorinated acrylic copolymer from Interpolymer Corporation
Syntran 1580	Acrylic copolymer from Interpolymer Corporation
Syntran DX302-1	Acrylic copolymer from Interpolymer Corporation
Onyxide 3300 33%	Miristalkonium saccharinate in alcohol from Stepan
Onyxide 3300	Miristalkonium saccharinate from Stepan
Stepanol WAC	Sodium lauryl sulfate from Stepan

Empicol 0335	Sodium alkyl sulfate from Hutsman
Petro 11 liquid	Sodium alkyl naphthalene sulfonate from Akzo
Eltesol SC40	Sodium cumene sulfonate from Hutsman
Genapol 26-L-5	Primary alcohol ethoxylate from Hoechst Celanese
Genapol 26-L-60	Primary alcohol ethoxylate from Hoechst Celanese
Kathon CG/ICP	Isothiazolinone from Rohm & Haas
Thymol Krist powder	Thymol from H. Reimer
Zonyl 7950	Fluorosurfactant from Dupont
Zelan 338	Polycarboxylate from Dupont
Dowanol PPH	1-phenoxy 2-propanol from Dow chem
Dowanol DPnP	Dipropylene glycol n-propyl ether from Dow chem
Isopropyl alcohol	Isopropyl alcohol
Fragrance	Proprietary fragrance from various suppliers

EXAMPLE RESULTS:

- The stain removal and carpet damage performances have been compared with
- 5 Resolve spot & stain carpet cleaner trigger and with Resolve pet stain & odour carpet cleaner trigger, considered as two of the market leaders in US for stain removal products.

Results for storage stability test:

10

In the table below are reported the storage rating values (SR) per each storage condition and the compatibility between onyxide 3300 and anionic species.

	Stability test rating (ready to use product)						Onyxide 3300 is compatible with the following anionics species:
Product	1 month			3 months			
	SR	SR	SR	SR	SR	SR	
	5°C	25°C	40°C	5°C	25°C	40°C	
Ex 1	2	1	1	2	1	1	Stepanol WAC, Syntran 1575
Ex 2	1	1	1	1	1	1	Stepanol WAC, Syntran 1575
Ex 3	0	1	1	1	1	3	Stepanol WAC, Syntran 1575
Ex 4	1	1	1	1	1	1	Stepanol WAC, Zelan 338, Zonyl 7950
Ex 5	0	1	1	1	1	5	Stepanol WAC, Zelan 338, Zonyl 7950
Ex 6	0	0	1	0	0	1	Stepanol WAC, Zelan 338, Zonyl 7950
Ex 7	0	0	2	5	2	6	Stepanol WAC, Zelan 338, Zonyl 7950
Ex 8	0	0	0	0	2	2	Stepanol WAC, Zelan 338, Zonyl 7950
Ex 9	0	0	0	0	0	2	Stepanol WAC, Zelan 338, Zonyl 7950
Ex 10	0	0	0	0	0	2	Stepanol WAC, Zelan 338, Zonyl 7950

	Stability test rating (dilutable product-manual and machine shampoos)			
Product	2 weeks			Onyde 3300 is compatible with the following anionics species:
	SR	SR	SR	
	5°C	25°C	40°C	
Ex 13	0	0	0	Eltesol SC40, Petro 11, Syntran 1580
Ex 14	0	0	0	Empicol 0335, Syntran DX 302-1
Ex 15	0	0	0	Empicol 0335, Syntran DX 302-1

5

Results for bactericidal efficacy test (ready to use product):

10

	Disinfectant efficacy					
	Staphylococcus aureus			Enterobacter aerogenes		
.	(bacteria reduction % with different contact times)					
	30 sec	1 min	5 min	30 sec	1 min	5 min
Reference: Resolve spot &	78.677	87.721	98.920	47.500	74.000	99.775
Ex 5	99.442	99.943	99.999	99.999	99.999	99.999
Ex 6	99.324	99.913	99.999	99.999	99.999	99.999

Ex 7	99.331	99.924	99.999	99.999	99.999	99.999
Ex 8	90.000	96.250	99.928	99.999	99.999	99.999
Ex 9	89.118	96.986	99.952	99.999	99.999	99.999
Ex 10	91.618	97.059	99.937	99.999	99.999	99.999

Results for stain removal tests (ready to use products):

- It has been used Resolve™ Spot and stain trigger and pet stain & odour trigger products as a references. 3 replicates has been considered per each product and the values reported in the table below are averaged.

	Stain removal rating					
	Mud	DMO	Grape juice	Coffee	Spaghetti sauce	Ink
Resolve™ Spot and stain trigger	3.7	3.3	3.0	3.1	3.2	4.2
Resolve™ pet stain & odour trigger	4.0	3.0	3.2	3.0	3.7	4.0
Ex 1	4.2	3.2	2.7	3.2	4.0	3.9
Ex 2	4.2	2.7	3.1	3.7	3.4	3.8
Ex 3	4.3	2.7	3.2	3.7	3.0	3.9
Ex 4	3.8	3.5	3.1	3.6	3.4	4.2
Ex 5	3.4	3.0	3.2	3.6	3.1	3.8

Results for carpet damage test and anti re-deposition (ready to use product):

It has been used again Resolve™ Spot and stain trigger and pet stain & odour trigger products as references. Per each product/carpet type has been considered 3 replicates and the values reported in the table below are averaged.

5

Product	Damage test				Anti re-depositio test
	Beige wool	Red wool	Blue nylon	Green polyester	
	rating	rating	rating	rating	%
Resolve™ Spot and stain trigger	0.7	0.6	0.1	0.1	10.3
Ex 4	0.6	0.6	0.5	0.0	8.0
Ex 5	1.0	0.8	0.2	0.0	2.5

Claims:

1. A detergent comprising:
 - a) 0.001% to 40% w/v of a bactericidally active water-insoluble cationic compound in its salt form wherein the counterion anion has at least one of the following properties:
 - 1) can generate a water-insoluble salt form, by water-insoluble we mean that less than 10% w/v dissolves in deionised water at 20°C, preferably less than 1%w/v;
 - 2) has a MW of less than 300, preferably less than 200 but greater than 50, preferably greater than 75;
 - 3) the dissociation constant (Kd) of the salt is less than 10^{-3} , preferably less than 10^{-6} ;
 - b) 0.001 to 40% w/v of an anionic surfactant; and
 - c) up to 98% w/v of water.
2. A detergent composition as claimed in claim 1 which additionally comprises 0.001 to 30% w/v of a non-ionic surfactant.
3. A detergent composition as claimed in claim 1 and 2 which additionally comprises 0.001 to 10% w/v of a superwetting agent.
4. A detergent composition as claimed in claim 3 wherein the super wetting agent is able to lower the surface tension of water to below 25 mN/m at concentrations of 0.0001 to 1% w/v.
5. A detergent composition as claimed in claims from 1 to 3 which additionally comprises 0.001% to 15% of a water-miscible organic solvent.


6. A detergent composition as claimed in claims 1 to 5 which additionally comprises 0.01-5%w/v of a chelating agent, 0.01-30%w/v of a polymer and up to 2%w/v of minor ingredients selected from perfumes, dyes, preservatives and antifoaming agents

5

7. A detergent composition described in claims 1 to 6 additionally comprises from 0.001 to 1%w/v of an additional bactericidally active product selected from essential oils (tea tree oil, citronella oil and thyme oil), phenols, alcohols, halogens, aldehydes and acids.

10


8. A detergent composition as claimed in any claim from 1 to 7 wherein the cationic compound is



wherein R1, R2, R3 and R4 are independently selected from alkyl, aryl or alkylaryl substituent of from 1 to 26 carbon atoms, and the entire cation portion of the molecule has a molecular weight of at least 165 and X is the counterion anion.

20

9. A detergent composition as claimed in claim 8 wherein the cationic compound is



wherein R2 and R3 are each independently the same or different C8 -C12 alkyl; or R2 is selected from C12-16 alkyl, C8-18 alkylethoxy or C8-18 alkylphenoxyethoxy and R3 is benzyl and X is the counterion anion.

30

10. A detergent composition as claimed in claim 7 or claim 8 wherein X^- is selected from saccharinate, alkyl and alkyl benzene sulfate, sulfonate and fatty acid.

5 11. Use of a bactericidally active water-soluble cationic compound in its salt form, wherein the counterion anion has at least one of the following properties:

1) can generate a water-insoluble salt form, by water-insoluble we mean that less than 10% w/v dissolves in deionised water at 20°C, preferably less than 1%w/v;

10

2) has a MW of less than 300, preferably less than 200 but greater than 50, preferably greater than 75;

15

3) the dissociation constant (K_d) of the salt is less than 10^{-3} , preferably less than 10^{-6} ;

as a bactericidally active component of a fabric treatment composition comprising from 0.001 to 40% w/v of an anionic surfactant.

20

THE PATENT OFFICE
22 AUG 2003
Received in Patents
International Unit

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.